

RESEARCH

Open Access



Non-deep physiological dormancy in seeds of *Euphorbia jolkinii* Boiss. native to Korea

Hye Jin Oh^{1†}, Un Seop Shin^{1,2†}, Seung Youn Lee^{1,3†}, Sang Yong Kim¹ and Mi Jin Jeong^{1*}

Abstract

Background: *Euphorbia jolkinii* Boiss. is a perennial species native to Jeju Island and the southern coastal area of Korea. Particularly on Jeju Island, the yellow flowers of *E. jolkinii* Boiss. have a high ornamental value because of their contrast with basalt. This study was conducted to investigate the effects of different temperatures (5, 15, 20, and 25 °C) and gibberellic acid (GA₃) concentrations (0, 10, 100, or 1000 mg/L) on seed dormancy and germination of *E. jolkinii*. In addition, we classified the seed dormancy type and compared types with those of other species in the same genus.

Results: The number of seeds with viable embryos and endosperms was approximately 66%. The final germination percentages at 5, 15, 20, and 25 °C were 51.7%, 83.5%, 2.6%, and 0.0%, respectively. In GA₃ concentration experiments, the final germination percentages of 0, 10, 100, and 1000 mg/L were 83.5%, 91.7%, 79.1%, and 83.4%, respectively, at 15 °C conditions, and 0.0%, 6.9%, 13.2%, and 27.3%, respectively, at 25 °C.

Conclusions: Germination improved at temperatures of 15 °C or lower. Furthermore, GA₃ treatment effectively reduced germination times. Thus, the seeds of *E. jolkinii* were classified as having non-deep physiological dormancy.

Keywords: Germination temperature, Gibberellic acid, Native plant, New ornamental plants, Physiological dormancy

Background

Euphorbia jolkinii Boiss. is a perennial herbaceous plant belonging to the Euphorbiaceae family (KPNI 2019). This plant is mainly distributed along the coasts of Jeju, Jeonnam, and Gyeongnam in Korea (Steinmann and Porter 2002). The genus *Euphorbia* has 1000–1600 species worldwide and is mainly distributed in subtropical and temperate zones (Steinmann and Porter 2002). Plants of the genus *Euphorbia* produce poisonous sticky white sap when damaged, and their flower structure is unique; therefore, they are grown for ornamental purposes or used for landscape decoration (*Euphorbia* PBI 2012). There are approximately 19 species native to Korea, including *E. jolkinii*, *E. esula* L., *E. helioscopia* L.,

and *E. humifusa* Willd. Ex Schltld (Park 2007; KNA 2017). *E. jolkinii* grows to 40–80 cm in height and forms a colony with more than 30 stems per population (Bae 2000; Shin et al. 2018a).

The inflorescence is cyathium, consisting of a number of stamens and one pistil, and yellow-green flowers bloom from March to May. The fruit is a capsule, there are bumps on the surface, and the seeds are round and flat, approximately 3 mm in diameter. Stems with green leaves grow in winter, and flowering begins in early spring and lasts for 3 months. In particular, the colorful yellow-green flowers contrast with basalt and have a high ornamental value in the native habitat on Jeju Island.

With the enactment of the Nagoya Protocol, biodiversity and biogenetic resources are highly valued, and efforts are being made to develop and industrialize native plants (Oh et al. 2018). *E. jolkinii* is a medicinal plant material, and leaves and stems have been used as

* Correspondence: mjeong121@korea.kr

[†]Hye Jin Oh, Un Seop Shin and Seung Youn Lee contributed equally to this work.

¹Division of Plant Resources, Korea National Arboretum, Yangpyeong 12519, Korea

Full list of author information is available at the end of the article



medicinal materials in oriental medicine. To verify and exploit these effects, studies on the pharmacological effects of extracts including their antioxidant and antibacterial activities, as well as melanin synthesis in melanoma cells have been conducted (Park et al. 2005; Kim et al. 2006; Kim et al. 2017). In addition, the unique, large flower structure and herbivorous form of *E. jolkinii* are considered to be of high ornamental value. To this end, researchers have evaluated the environment and growth characteristics of native habitats (Shin et al. 2018a), the vase life of cut flowers (Oh et al. 2018; Song et al. 2019a), cuttings for propagation (Shin et al. 2017), and cultivation according to pot size and nutrient conditions (Shin et al. 2018b). However, the dormancy characteristics of seeds and the conditions and characteristics of seed germination have not yet been reported.

Seed dormancy refers to the plant's self-suppression of germination under adverse environmental conditions for survival and is classified as physiological dormancy (PD), morphological dormancy (MD), morphophysiological dormancy (MPD), physical dormancy (PY), and combinational dormancy (PY + PD) (Baskin and Baskin 1998; Baskin and Baskin 2004; Fenner and Thompson 2005). In PD, dormancy is affected by cold stratification, after-ripening, and gibberellic acid (GA) treatment. PD can be divided into three types: non-deep PD, intermediate PD, and deep PD, depending on the depth of dormancy (Nikolaeva 1977; Baskin and Baskin 2014). Non-deep PD is the most common form of seed dormancy. This type of dormancy can be broken by short periods of chilling stratification, after-ripening (dry storage), or gibberellin treatment. Intermediate PD usually requires at least 2 months of chilling or gibberellin application as a substitute for chilling, whereas deep PD requires a long period of at least 3 months of chilling stratification to relieve dormancy and generally does not respond to exogenous gibberellin application (Geneve 2003; Baskin and Baskin 2004). MD and MPD are ecophysiological characteristics that occur because seeds have immature embryos at the time they are separated from the parent plant (Adams et al. 2005a, 2005b). Immature embryos are defined not by the size of the embryo, but by how much the embryo must still grow before germination (or before the seed coat is split) (Baskin and Baskin 2005; Chen et al. 2015). Seeds in the genus *Euphorbia* have been reported to be mostly nondormant or to show PD (Best et al. 1980; Sen and Chatterji 1966; Gómez and Espadaler 1997; Narbona 2002). *E. esula* can break dormancy after a 3-month period of after-ripening treatment (Best et al. 1980), and *E. caducifolia*, *E. characias*, and *E. nicaensis*, which are not native to Korea, do not exhibit dormancy. However, none of these studies have clearly revealed the morphology and development of the embryo. Therefore, detailed

research is needed to determine the characteristics of non-dormancy and PD.

Accordingly, in this study, we aimed to obtain basic data for the development of new ornamental plants by identifying optimal germination conditions and classifying dormancy types for *E. jolkinii* seeds.

Materials and methods

Plant material

The *E. jolkinii* seeds used in this experiment were collected on June 14, 2018, in Seongsan-eup, Seogwipo-si, Jeju (33° 18' 23.0", 126° 49' 36.0"). The collected seeds were dried at room temperature (approximately 25 °C) for 2 weeks in the laboratory and then used in experiments.

Morphological characteristics of seeds

To observe the morphology of the seeds, 20 seeds were randomly selected. The seeds were cut in half with a double-edged razor (stainless blade; Dorco, Seoul, Korea). The sections were photographed at 50–60× magnification using a Dino-Life Edge Digital Microscope (AM 3111 Dino-Lite Premier; AnMo Electronics Co., Taiwan), and the secession, internal and external shapes of the seeds, and germination process were observed.

Water uptake test

To investigate whether seeds showed PY, the water uptake rate was checked in three replicates of 30 seeds. Seeds were placed in plastic Petri dishes (90 mm × 10 mm) on top of two filter papers (Whatman No.1; GE Healthcare, Buckinghamshire, UK) moistened with distilled water at room temperature. Seeds were weighed after 0, 3, 6, 9, 12, 24, 48, 72, and 96 h of incubation. Percent water uptake was calculated as $\%W_s = \frac{[W_h - W_i]}{W_i} \times 100$, where W_s is the increase in mass of the seed, W_h is the seed mass after a given interval of imbibition, and W_i is the initial seed mass at 0 h.

Temperature treatments

Seeds were soaked in a solution of 1000 mg/L benomyl (Dongbu Benomyl; Farm Hannong, Seoul, Korea) for 1 h. After sterilization, the samples were washed with distilled water five times. Seeds were placed in plastic Petri dishes (90 mm × 10 mm) on top of two filter papers (Whatman No.1; GE Healthcare) moistened with distilled water. To determine the optimum temperature for germination, seeds were incubated at 5, 15, 20 and 25 °C with light (16 h, cool white fluorescent lamp, approximately 25 $\mu\text{mol}/\text{m}^2/\text{s}$) or darkness in a constant temperature growth chamber (Multi-Room Incubator; WiseCube, Wonju, Korea).

GA₃ treatments

To promote germination, seeds were soaked in GA₃ solutions of different concentrations (0, 10, 100, and 1000 mg/L) for 24 h at room temperature. Then, seeds were incubated at 15 °C and 25 °C under light conditions of 16 h light/8 h dark. All treatments were conducted in triplicates with 30 seeds. Germination was checked at day 3 of the experiment. In growth chambers, the photosynthetic photon flux was approximately 25 μmol/m²/s during the light period, and a 16-h light/8-h dark photoperiod was used. Sterile water was periodically administered to keep the filter paper moist.

Cold stratification treatments

Sterilized seeds were treated for 0, 4, 8, and 12 weeks at 4 °C and then incubated at 25 °C. The experiment was performed in three repetitions of 30 seeds, and germination was measured at 1-week intervals.

After-ripening treatments

The harvested seeds were dried and stored at room temperature for 0, 3, and 6 weeks. At the end of the treatment period, the seeds were sterilized and incubated at 25 °C. The experiment was performed in three repetitions of 30 seeds, and germination was measured at 1-week intervals.

Germination investigation

Germination was defined as the point at which radicles emerged at least 1 mm beyond the seed coat. Radicle emergence was checked every 3 days. After the final germination observation, poor-quality seeds were excluded from calculation of the germination percentage. The final germination was measured after 16 weeks of seed incubation. The germination characteristics, final germination percentage (FGP), day of germination, mean germination time (MGT) and days to 50% germination (T_{50}) in days were calculated, as follows:

$$\text{FGP} = N_g/N_t \times 100$$

where N_g is the number of final germinated seeds and N_t is the total number of seeds tested (Al-Mudaris 1998; Gharineh et al. 2004).

$$T_{50} = T_i + [(N + 1)/2 - N_i]/(N_j - N_i) \times (T_j - T_i)$$

where N is the final number of seeds germinating, and N_i and N_j are the total numbers of seeds germinated by adjacent counts at times T_i and T_j (Coolbear et al. 1984).

$$\text{MGT} = \Sigma (N_i \times T_i)/N$$

where N_i is the number of seeds germinated, T_i is the number of days from the start of the experiment, N is

the total number of germinated seeds at the end of the experiment (Naylor 1981).

Data analysis

Data were subjected to statistical analysis using SAS 9.4 software (SAS Inst. Inc., Cary, NC, USA). Statistical evaluation was conducted using analysis of variance, and significant data were analyzed further using Tukey's honestly significant difference test ($p \leq 0.05$).

Results

Morphological characteristics of seeds

E. jolkinii seeds are surrounded by a hard pericarp with bumps (KBIS 2019). The seeds were dark brown and globose or spherical (Fig. 1A). The surface appeared smooth, but had a mesh structure (areolate) (KNA 2017). The collected seeds were 2.12–2.85 mm in length and 2.10–2.35 mm in width, with an average of 2.18 mm. The weight of 100 seeds was approximately 473 mg, and the percentage of sound seeds was approximately 66%. After cutting the seeds, no endosperm vigor was observed in some of the seeds (Fig. 1D).

Water uptake rate

The weight of the seeds increased by 21 % compared to the initial dry weight after 6 h of water absorption and by 39% after 24 h of water absorption (Fig. 2). After showing a 50% increase in water uptake, the seeds were saturated and did not increase further in weight after 72 h.

Temperature control of seed germination

The FGPs at 5, 15, 20, and 25 °C were 51.7%, 83.5%, 2.6%, and 0.0%, respectively (Fig. 3). No seeds germinated under the four temperatures for four weeks. Less than 5% of the seeds germinated at 20 °C and 25 °C. At 5 °C and 15 °C, germination started after 47.3 and 39.0 days, respectively (Table 1). The mean germination times at 5 °C and 15 °C were 53.8 and 82.3 days, respectively. The FGP was higher at 15 °C, but the germination progression was faster at 5 °C. In addition, T_{50} values were 48.7 and 85.0 days at 5 and 15 °C, respectively.

Effects of GA₃ treatment on seed germination

GA₃ effectively breaks seed dormancy, promotes embryo development, and enhances germination (Finch-Savage 2006; Baskin and Baskin 2014; Cho and Lee 2017). GA₃ treatment affected the germination of *E. jolkinii* seeds at 15 °C and 25 °C; in particular, the germination was promoted effectively at 15 °C. Following treatments with 0, 10, 100, or 1000 mg/L of GA₃, the germination percentages were 83.5%, 91.7%, 79.1%, and 83.4%, respectively, at 15 °C and 0%, 6.9%, 13.2%, and 27.3%, respectively, at 25 °C (Fig. 4). There were significant differences in the day of germination according to GA₃ concentration




Fig. 1 Germination process of *Euphorbia jolkinii* Boiss. seeds. **A** A fruit and a seed. **B** Inner morphology. **C** A radicle. **D** A festered seed. **E, F** Seedling emergence. **G, H** Flowering of *E. jolkinii* in a native habitat. en, endosperm; em, embryo. Scale bars = 1 mm




Fig. 2 Water uptake by seeds of *Euphorbia jolkinii* Boiss. as represented by an increase in mass. Seeds were incubated at room temperature for 96 h. Error bars indicate means \pm standard errors of three replicates of 30 seeds




Fig. 3 Germination of *Euphorbia jolkinii* Boiss. seeds as affected by different temperature treatments. Error bars indicate means ± standard errors of three replicates

(Table 2). At 15 °C, the GA₃ control started to germinate 39 days after the start of the experiment, whereas seeds treated with 100 and 1000 mg/L GA₃ started germination on days 13 and 7, respectively. The mean times to germination after 0, 10, 100, and 1000 mg/L GA₃ treatment were 82.3, 84.7, 34.8 days, and 34.8 days, respectively, and the T₅₀ values were 85.0, 95.0, 47.7, and 41.3 days, respectively (Table 2).

Cold stratification treatments

To assess the dormancy of *E. jolkinii* the germination percentage was observed for 9 weeks after cold stratification treatment for 4, 8, and 12 weeks. However, the germination percentage was less than 10% irrespective of the treatment duration.

After-ripening treatments

Germination was observed for 17 weeks after 3 and 6 months of after-ripening treatment at room temperature; however, the seeds did not germinate.

Discussion


The seeds were divided into basal, peripheral, and axile, according to the shape of the embryo (Martin 1946). In the basal type, the embryo does not exceed half the size of the seed and is located at the lower end. The peripheral type has a large embryo and is curved along the seed coat. In the axile type, the embryo varies and is located in the center of the seed (Song et al. 2019b). *E. jolkinii* seeds were axile, with the embryo located along the center of the seed (Fig. 1B). Seeds of plants native to the temperate regions are often separated from the parent without developing enough embryos, and seeds with such immature embryos are MD- or MPD-type seeds with a ratio of 0.5% or less. We confirmed that the length of the embryo in *E. jolkinii* seeds accounted for approximately 80–90% of the length of the seed, and the radicles protruded without any difference in the length of the embryo just before germination (Fig. 1C). Therefore, there was no MD in *E. jolkinii* seeds.

If germination does not occur because of the impermeability of the seed coat or pericarp, the seed is classified as showing PY (Baskin et al. 2000). If the water

Table 1 Effect of temperatures on the germination of *Euphorbia jolkinii* seeds

Temperature (°C)	Days to germination	Mean times to germination	T ₅₀
5	47.3 ± 1.3b ²	53.8 ± 1.8b	48.7 ± 1.3b
15	39.0 ± 4.0b	82.3 ± 2.9a	85.0 ± 3.0a
20	102.0 ± 0.0a	–	–
25	–	–	–
P-value	0.0012	0.0011	0.0004

²The different letters are significantly different (p ≤ 0.05) using Tukey's HSD test (n = 3)



uptake rate is more than 20% within 24 h, the seed is judged to have water permeability (Baskin and Baskin 2003). Therefore, we found that there was no PY in *E. jolkinii* seeds, which had increased water absorption within 6 h.

E. jolkinii had the highest germination percentage at 15 °C. In comparison, *E. boetica* had a germination percentage of 73% at 17/21 °C (night/day, 13 h/11 h) and a T_{50} of 7.4 days, and *E. heterophylla* had a germination percentage of about 80% at 25 °C and 30 °C (Cecilia and Jarbas 2000; Narbona et al. 2007). *E. boetica* grows in lowland pine forests or acidic soils (Benedí et al. 1997), and *E. heterophylla* grows as weeds in tropical and subtropical America (Hutchinson and Dalziel 1958; Wilson 1981). Different plants have different germination conditions, and even similar plants may show major differences in the germination characteristics when their native habitats differ at the national level (Baskin et al. 1989). Thus, we confirmed that the ideal environmental conditions for the germination of *E. jolkinii*, which is distributed only along the southern coast of Korea, were different from other species of the genus *Euphorbia*.

In non-deep PD, dormancy of seeds can be broken by various treatments, including 1–2 months of cold

stratification treatment, after-ripening, and GA treatment. In intermediate PD, dormancy breaking through GA treatment is limited to some species, and it is possible to break dormancy through cold stratification treatment for 2–3 months (Geneve 2003; Baskin and Baskin 2004). In addition, it is possible to partially satisfy the low-temperature requirement for breaking dormancy through dry storage such as post-aging. In deep PD, the dormancy of seeds cannot be broken by GA treatment, but it is possible to break dormancy through 3–4 months of low-temperature or high-temperature wet treatment (Geneve 2003; Baskin and Baskin 2004). In this study, we showed that GA_3 treatment could effectively improve the germination percentages of *E. jolkinii* seeds. The germination percentage was 79.1–83.4% at 100 and 1000 mg/L, but there were no significant differences between these two concentrations. In addition, the day of germination, mean time to germination, and T_{50} were shortened in the presence of 100 and 1000 mg/L GA_3 . Indeed, at these concentrations, the day to germination was 3–5 times earlier, the mean time to germination was shortened 2 times, and the T_{50} was shortened by 2.0–2.3 times compared with the control. Therefore, GA_3 treatment improved the germination percentage

Table 2 Effect of GA_3 concentrations on the germination of *Euphorbia jolkinii* seeds at 15 °C

GA_3 concentration (mg/L)	Days to germination	Mean times to germination	T_{50}
0	39.0 ± 4.0b ^z	82.3 ± 2.9a	85.0 ± 3.0a
10	83.5 ± 1.5a	84.7 ± 2.4a	95.0 ± 0.0a
100	13.0 ± 1.0c	34.8 ± 2.8b	47.7 ± 6.2b
1000	7.0 ± 0.0c	34.8 ± 9.1b	41.3 ± 8.4b
P-value	0.0001	0.0004	0.0011

^zThe different letters are significantly different ($p \leq 0.05$) using Tukey's HSD test ($n = 3$)

and shortened the mean time to germination, confirming that the seeds showed non-deep PD.

Among native plants in Korea, gibberellin treatment has been shown to improve or promote seed germination. Germination percentages of *Primula modesta* var. *hannasanensis* T.Yamaz. were 95.0% and 95.5% at GA₃ concentrations of 200 and 500 mg/L, respectively, which were 56.0–69.0% higher than those of the control treatment (Cho and Lee 2017). Additionally, the germination percentage (72.5%) of *Tiarella polyphylla* D. Don was improved by GA₃ treatment at a concentration of 1000 mg/L compared with the control treatment (33.3%) (Choi et al., 2018). Moreover, the germination percentages of *Veronica kiusiana* var. *diamantiaca* (Nakai) T. Yamaz. were enhanced by GA₃ treatment to 71.3%, 87.2%, and 91.7% at GA₃ concentrations of 50, 100, and 1000 mg/L, respectively, compared with that of the control (44.7%) (Song et al. 2019b). *Zanthoxylum piperitum* (Kim et al. 1997), *Exochorda serratifolia* (Lee et al. 2006), *Pinus pumila* (Lim et al. 2015), *Lychnis wilfordii* (Regel) Maxim (Ryu et al. 2017), and many other species have also been shown to have improved germination percentages following gibberellin treatment, suggesting that this treatment breaks the PD of seeds. Accordingly, these findings showed that the concentration and time of gibberellin treatment were critical factors for improving germination percentages.

Conclusion

Wild native species have a wide distribution range, and it is not easy to establish an appropriate environment for seed germination. *E. jolkinii* Boiss. seeds require a longer time to germinate than other species in the same genus *Euphorbia*. In this study, we confirmed that these seeds showed no MD or PY, as demonstrated by embryo observation and water permeability. When germination was observed at various temperatures, we found that the optimum temperature for germination was 15 °C, although the mean germination time at 15 °C was very late (82.3 days), and PD was confirmed. Additionally, the days to germination, mean time to germination, and T₅₀ were shortened following GA₃ treatment at concentrations of 100 and 1000 mg/L, thus classifying *E. jolkinii* seeds as showing non-deep PD. Overall, we confirmed that the germination of *E. jolkinii* seeds was improved at temperatures below 15°C. In addition, cold stratification and after-ripening, GA₃ individual treatments were not effective for germination. However, treatment with GA₃ at 15°C reduced the mean germination time. Scientific conclusions on these environmental conditions and dormancy will be useful for establishing a mass production system for commercial use and ecological management.

Abbreviations

PD: Physiological dormancy; MD: Morphological dormancy; MPD: Morphophysiological dormancy; PY: Physical dormancy; FGP: Final germination percentage; T₅₀: Days to 50% germination; MGT: Mean germination time

Acknowledgements

Not applicable.

Authors' contributions

OHJ and SUS carried out the study, analyzed the data, and wrote the manuscript. LSY conceived the study design, analyzed the data, and reviewed the manuscript. KSY collected and analyzed the data. JMJ participated in the study and reviewed and edited the manuscript. All authors read and approved the final manuscript.

Funding

This study was funded by the project "Development of Year-round Cultivation and Flowering Control Technique for Native Wild Flowers Commercialization and its Diversification of Utilization (KNA1-2-33, 17-8)" funded by the Korea National Arboretum.

Availability of data and materials

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Division of Plant Resources, Korea National Arboretum, Yangpyeong 12519, Korea. ²Seed Viability Research Team, Baekdudaegan National Arboretum, Bongwa 36209, Korea. ³Division of Horticulture & Medicinal Plant, Andong National University, Andong 36729, Korea.

Received: 7 September 2021 Accepted: 30 September 2021

Published online: 19 October 2021

References

- Adams CA, Baskin JM, Baskin CC. Trait stasis versus adaptation in disjunct relict species: evolutionary changes in seed dormancy-breaking and germination requirements in a subclade of *Aristolochia* subgenus *Siphisia* (Piperales). *Seed Sci Res.* 2005a;15(2):161–73. <https://doi.org/10.1079/SSR2005207>.
- Adams CA, Baskin JM, Baskin CC. Comparative morphology of seeds of four closely related species of *Aristolochia* subgenus *Siphisia* (Aristolochiaceae, Piperales). *Bot J Linnean Soc.* 2005b;148(4):433–6. <https://doi.org/10.1111/j.1095-8339.2005.00402.x>.
- Al-Mudaris M. Notes on various parameters recording the speed of seed germination. *Tropenlandwirt.* 1998;99:147–54.
- Bae KH. Medicinal plants in Korea. Seoul; Kyohaksa. 2000.
- Baskin CC, Baskin JM. Seeds: ecology, biogeography, and evolution of dormancy and germination. San Diego: Academic Press; 1998.
- Baskin CC, Baskin JM. When breaking seed dormancy is a problem: try a move-along experiment. *Nativ Plant J.* 2003;4(1):17–21. <https://doi.org/10.3368/npj.4.1.17>.
- Baskin CC, Baskin JM. Underdeveloped embryos in dwarf seeds and implications for assignment to dormancy class. *Seed Sci Res.* 2005;15(4):357–60. <https://doi.org/10.1079/SSR2005224>.
- Baskin CC, Baskin JM. Seeds: ecology, biogeography, evolution of dormancy and germination. 2nd ed. San Diego: Elsevier; 2014.
- Baskin JM, Baskin CC. A classification system for seed dormancy. *Seed Sci Res.* 2004;14(1):1–16. <https://doi.org/10.1079/SSR2003150>.

- Baskin JM, Baskin CC, Li X. Taxonomy, anatomy and evolution of physical dormancy in seeds. *Plant Spec Biol.* 2000;15(2):139–52. <https://doi.org/10.1046/j.1442-1984.2000.00034.x>.
- Baskin JM, Baskin CC, Spooner DM. Role of temperature, light and date seeds were exhumed from soil on germination of four wetland perennials. *Aquat Bot.* 1989;35(3-4):387–94. [https://doi.org/10.1016/0304-3770\(89\)90009-0](https://doi.org/10.1016/0304-3770(89)90009-0).
- Benedí C, Molero J, Simón J, Vicens J. Euphorbia. In: Castroviejo S, Aedo C, Benedí C, Lainz MR, Muñoz-Garmendia F, Nieto-Feliner G, Paiva J, editors. *Flora Iberica*, vol. 7. Madrid: Real Jardín Botánico de Madrid-CSIC; 1997. p. 210–85.
- Best KF, Bowes GG, Thomas AG, Maw MG. The biology of Canadian weeds. 39. *Euphorbia esula* L. *Can J Plant Sci.* 1980;60(2):651–63. <https://doi.org/10.4141/cjps80-092>.
- Cecilia NKS, Jarbas FG. Seed reserve composition and mobilization during germination and initial seedling development of *euphorbia geterophylla*. *Rev Bras Fisiol Veg.* 2000;12(3):226–46. <https://doi.org/10.1590/S0103-313200000300006>.
- Chen SY, Tsai YH, Baskin CC, Baskin JM, Chien CT. Seed dormancy and germination of the three tropical medicinal species *Gomphandra luzoniensis* (Stemonuraceae), *Nothapodytes nimmoniana* (Icacinaeae) and *Goniothalamus amuyon* (Annonaceae). *Seed Sci Res.* 2015;25(01):57–63. <https://doi.org/10.1017/S0960258514000385>.
- Cho JS, Lee CH. Dormancy and germination characteristics of alpine modest primrose (*Primula modesta* var. *hannasanensis* T.Yamaz.) seeds. *Korean J Plant Res.* 2017;30:372–7.
- Choi H, Lee SY, Rhie YH, Lee JH, Kim SY, Lee KC. Seed dormancy type and germination characteristics in *Tiarella polyphylla* D. Don native to Korea. *Korean J Plant Res.* 2018;31:363–71.
- Coolbear P, Francis A, Grierson D. The effect of low temperature pre-sowing treatment under the germination performance and membrane integrity of artificially aged tomato seeds. *J Exp Bot.* 1984;35:1609–17.
- Euphorbia Planetary Biodiversity Inventory (PBI) Project. 2012. <http://www.euphorbiaceae.org/index.html>.
- Fenner M, Thompson K. The ecology of seeds. NY: Cambridge University Press; 2005. <https://doi.org/10.1017/CBO9780511614101>.
- Finch-Savage WE, Leubner-Metzger G. Seed dormancy and the control of germination. *New Phytol.* 2006;71(3):501–23. <https://doi.org/10.1111/j.1469-8137.2006.01787.x>.
- Geneve RL. Impact of temperature on seed dormancy. *Hortscience.* 2003;38(3):336–41. <https://doi.org/10.21273/HORTSCI.38.3.336>.
- Gharineh MH, Bakhshandeh A, Ghasemi-Golezani K. Vigor and seed germination of wheat cultivar in Khuzestan environmental condition. *Sci J Agric.* 2004;27:65–76.
- Gómez C, Espadaler X. Manipulación por hormigas de Semillas de *Euphorbia characias* (Euphorbiaceae) dentro del hormiguero. *Sci Gerund.* 1997;23:53–61.
- Hutchinson J, Dalziel JM. Flora of west tropical Africa, Vol 1 (ii) (2nd edition, revised by Keay RWJ, 1973). London: Crown Agents for Overseas Governments and Administrations; 1958. p. 297–828.
- Kim JY, Lee JA, Yoon WJ, Oh DJ, Jung YH, Lee WJ, et al. Antioxidative and antimicrobial activities of *euphorbia jolkinii* extracts. *Korean J Food Sci Technol.* 2006;38:699–706.
- Kim MJ, Kim SY, Hyun KH, Kim DS, Kim SY, Hyun CG. Effects of *Rumex acetosella*, *Sonchus oleraceus* and *euphorbia jolkinii* extracts on melanin synthesis in melanoma cells. *Korean Soc Biotechnol Bioeng J.* 2017;32:187–92.
- Kim SJ, Shin JH, Kim KJ, Park SD, Choi BS, Kim KU. Effect of GA₃, kinetin and physical treatment on the seed germination of *Zanthoxylum piperitum* a.P. DC. *Korean J Med Crop Sci.* 1997;5:43–8.
- Korea Biodiversity Information System (KBIS). 2019. <http://www.nature.go.kr/kbi/plant/pilbk/selectPlantPilbkDtl.do>.
- Korea National Arboretum. Seed atlas of Korea. Seoul: Sumeunki Publishing Co.; 2017.
- Korean Plant Names Index (KPNI). 2019. <http://www.nature.go.kr/kpni/index.do>.
- Lee HD, Kim SD, Kim HH, Lee JW, Kim JH, Lee CH, Lee CH. Effects of storage method, growth regulator, and inorganic salt on the seed germination of *Eoachorda serratifolia* S. Moore. *Korean J Hort Sci Technol.* 2006;24:90–4.
- Lim HI, Kim GN, Jang KH, Park WG. Effect of wet cold and gibberellin treatments on germination of dwarf stone pine seeds. *Korean J Plant Res.* 2015;28(2):253–8. <https://doi.org/10.7732/kjpr.2015.28.2.253>.
- Martin AC. The comparative internal morphology of seeds. *Am Midl Nat.* 1946;36(3):513–660. <https://doi.org/10.2307/2421457>.
- Narbona E. Estrategias reproductivas de dos especies perennes de euphorbia. Ph. D thesis. Sevilla: Universidad de; 2002.
- Narbona E, Arista M, Ortiz PL. Seed germination ecology of the perennial *euphorbia boetica*, an endemic spurge of the southern Iberian Peninsula. *Ann Bot Fennici.* 2007;44:276–82.
- Naylor REL. An evaluation of various germination indices for predicting differences in seed vigour in Italian ryegrass. *Seed Sci Technol.* 1981;9:593–600.
- Nikolaeva MG. Factors controlling the seed dormancy pattern. In: Khan AA, editor. *The physiology and biochemistry of seed dormancy and germination*. Amsterdam: North-Holland; 1977. p. 51–74.
- Oh HJ, Shin US, Lee JS, Kim SY, Lee SY. Evaluation of postharvest cut flower characteristics of a new ornamental crop known as *euphorbia jolkinii* Boiss. *Flower Res J.* 2018;26(3):132–41. <https://doi.org/10.11623/frj.2018.26.3.07>.
- Park KR. Euphorbiaceae. In: the genera of vascular plants of Korea. Flora of Korea editorial committee. Seoul: Academy Publishing Co.; 2007.
- Park SH, Kim JA, Hua XG, Lee CG, Choi JY. Isolation of 5α-reductase inhibitors from *euphorbia jolkinii*. *Korean J Pharmacogn.* 2005;36:9–16.
- Ryu SH, Rhie YH, Lee SH, Ko CH, Lee JH, Lee KC. Effect of after-ripening, cold stratification, and GA₃ treatment on *Lychnis wilfordii* (regel) maxim seed germination. *Korean J Hort Sci Technol.* 2017;35:525–33.
- Sen DN, Chatterji UN. Eco-physiological observations of *euphorbia caducifolia* Haines. *Sci Cult.* 1966;32:317–9.
- Shin US, Lee JS, Oh HJ, Kim SY, Lee SY. Population distribution and growth environment of *euphorbia jolkinii* Boiss. In Jeju Island, Korea. *Flower Res J.* 2018a;26(3):109–16. <https://doi.org/10.11623/frj.2018.26.3.04>.
- Shin US, Oh HJ, Lee JS, Min SY, Kim SY, Lee SY. Vegetative growth responses of *euphorbia jolkinii* Boiss. As affected by pot size and nutrients treatment. *Hortic Sci Technol.* 2018b;36:189–90.
- Shin US, Oh HJ, Min SY, Lee JS, Lee KC, Kim SY, et al. Effect of cutting date, photoperiod and plant growth regulators on rooting of *euphorbia jolkinii* Boiss. Stem cuttings. *Hortic Sci Technol.* 2017;35:189–90.
- Song SJ, Park HB, Kim JS, Oh HJ, Kim SY, Jeong MJ, et al. Effect of holding solution on vase life of a new ornamental crop known as *euphorbia jolkinii* Boiss. *Korean J Plant Res.* 2019a;32:312–7.
- Song SJ, Shin US, Oh HJ, Kim SY, Lee SY. Seed germination responses and interspecific variations to different incubation temperature in eight Veroniceae species native to Korea. *Korean J Hort Sci Technol.* 2019b;37(1):20–31. <https://doi.org/10.12972/kjhst.20190003>.
- Steinmann VW, Porter JM. Phylogenetic relationships in Euphorbiaceae (Euphorbiaceae) based on ITS and *ndhF* sequence data. *Ann Mo Bot Gard.* 2002;89:53–490.
- Wilson AK. *Euphorbia heterophylla*: a review of distribution, importance and control. *Trop Pest Manag.* 1981;27(1):32–8. <https://doi.org/10.1080/09670878109414169>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

